

Implementation of Enhanced Recovery After Surgery (ERAS) protocol in off-pump coronary artery bypass graft surgery. A prospective cohort feasibility study

Michał Borys¹, Sławomir Żurek², Arkadiusz Kurowicki², Beata Horeczy³, Bartłomiej Bielina³, Justyna Sejboth⁴, Bogumiła Wołoszczuk-Gębicka³, Mirosław Czuczwar¹, Kazimierz Widenka²

¹Second Department of Anesthesia and Intensive Care, Medical University of Lublin, Lublin, Poland

²Clinical Department of Cardiac Surgery, Medical Faculty, University of Rzeszów, Rzeszów, Poland

³Department of Anesthesiology and Intensive Care with the Center for Acute Poisoning, St. Jadwiga Provincial Clinical Hospital, Rzeszów, Poland

⁴Department of Anaesthesia and Intensive Therapy, School of Health Sciences in Katowice, Medical University of Silesia in Katowice, Poland

Abstract

Background: Coronary artery bypass graft (CABG) is the most commonly performed cardiac surgery procedure. Although some complications related to the cardiopulmonary bypass circuit are avoided during off-pump CABG (OP-CABG) procedures, prolonged mechanical ventilation and severe postoperative pain are still important issues.

Methods: This prospective cohort study aimed to assess the impact of the institutional Enhanced Recovery After Surgery (ERAS) protocol in patients undergoing OP-CABG. This protocol contained several modifications to the perioperative period, among which bilateral erector spinae plane block, remifentanyl infusion, and patient-controlled analgesia (PCA) with oxycodone were the most important factors (ERAS group). The ERAS group was compared with the retrospective cohort (same surgeon) before the ERAS protocol was implemented (standard care group). The outcomes measured included the postoperative mechanical ventilation time, ICU and hospital stay, postoperative drainage time, postoperative troponin T level, pain severity evaluated via a numerical rating scale, and the total consumption of opioids in both groups of patients.

Results: Overall, 57 patients were analyzed – 29 in the ERAS group and 28 in the standard care group. The time of mechanical ventilation, thoracic drainage, and ICU and hospital stay was shorter in the ERAS group than in the standard care group. The pain was less intense in the ERAS patients, and the postoperative opioid demand was reduced. Moreover, the increase of the postoperative troponin T concentration was lower in the ERAS group.

Conclusions: Our study showed that ERAS protocol implementation could improve patient outcomes after OP-CABG surgery.

Key words: patient-controlled analgesia, coronary artery bypass graft, Enhanced Recovery After Surgery, erector spinae plane block, numerical rating scale.

Anestezjologia Intensywna Terapia
2020; 52, 1: 10–14

Otrzymano: 29.08.2019,
zaakceptowano: 06.01.2020

ADRES DO KORESPONDENCJI:

Dr. Michał Borys, Second Department of Anesthesia and Intensive Care, Medical University of Lublin, 16 Staszica Str., 20-081 Lublin, Poland, e-mail: michalborys1@gmail.com

Coronary artery bypass graft (CABG) is the most commonly performed cardiac surgery procedure [1]. Approximately 62 procedures are performed per 100,000 inhabitants in Europe each year [2]. However, the complication rate is still relatively high, even after scheduled CABG [3].

Among the many problems related to CABG, the most important aspect, which might be affected by anesthetic care, is postoperative pain [4]. Many patients still suffer from severe pain after CABG

procedures [5]. Regional anesthetic techniques are effective in pain alleviation and are associated with decreased mechanical ventilation time [6]. Still, most cardiac surgery patients are mechanically ventilated under sedation during the postoperative period [4]. Prolonged mechanical ventilation is associated with postoperative respiratory complications [7].

Effective pain treatment is one of the mainstays of the Enhanced Recovery After Surgery (ERAS) protocol [8]. However, this strategy involves a simulta-

neous modification of many aspects of perioperative care.

Our previous study showed that the implementation of the ERAS protocol was possible in patients undergoing mitral/tricuspid valve repair via the right mini-thoracotomy [9]. For the mini-thoracotomy, the institutional ERAS protocol consisted of patient education, avoidance of prolonged pre-op fasting, and pharmacological premedication. Moreover, in this study, we used a unilateral erector-spinae-plane (ESP) block before the operation and continuous infusion of remifentanyl during the surgery and for two hours during the early postoperative period, followed by patient-controlled analgesia with intravenous oxycodone. The ESP block provided efficient pain control, enabling a significant reduction in the perioperative opioid dosage, early extubation, and a shorter patient stay in the ICU. In the conclusion of our previous study, we stated that the usefulness of ESP block in other types of procedures needs to be studied more extensively.

Our current study aimed to investigate whether a preemptively performed, bilateral ESP block combined with intraoperative remifentanyl infusion could be useful and safe in patients undergoing off-pump CABG (OP-CABG). These patients were compared with a retrospective cohort that was anesthetized with fentanyl, but without any regional analgesia techniques.

The main goal of the study was to compare mechanical ventilation time in the two groups of patients. The other outcomes included ICU and hospital stay, postoperative drainage time, postoperative troponin T level, pain severity, total opioid consumption, and perioperative complications in both patient groups.

METHODS

This was a prospective cohort feasibility study conducted in a tertiary cardiac surgery department. Before patient recruitment, the study protocol was approved by the Bioethics Committee of the Medical University of Lublin in Lublin, Poland (permit number KE-0254/219/2018). Informed consent was obtained from each patient, and the study was conducted in accordance with the tenets of the Declaration of Helsinki for medical research involving human subjects.

Participants

The patients subsequently undergoing OP-CABG had perioperative treatment according to the institutional ERAS cardio protocol (ERAS group). All patients signed the written informed consent form. Their perioperative course was compared with the retrospective data from patients who had the same

surgery performed by the same surgeon, but before the ERAS protocol implementation (standard care group).

Intervention

In the ERAS group prior to the induction of general anesthesia, a single-shot ESP block was performed bilaterally as described in our previous study [9], and 0.2 mL kg⁻¹ of 0.375% ropivacaine (Ropimol, Molteni, Italy) was administered on each side. The total volume of local anesthetic solution did not exceed 40 mL.

Induction of general anesthesia was provided with 0.2–0.4 mg kg⁻¹ of etomidate (Hypnomidate, Janssen-Cilag International NV, Belgium), 0.4–0.6 µg kg⁻¹ of remifentanyl (Ultiva, GlaxoSmithKline, UK), and 0.6 mg kg⁻¹ of rocuronium (Esmeron, NV Organon, The Netherlands). Sevoflurane 0.5 MAC (age-adjusted, Sevorane, Abbvie, USA), remifentanyl infusion according to the target control protocol (4–8 ng mL⁻¹), and incremental doses of rocuronium were used for maintenance.

Patients received an intravenous bolus of oxycodone (0.1 mg kg⁻¹) about 30 minutes before the end of the surgery. Remifentanyl infusion was decreased to the target plasma concentration of 0.5–2 ng mL⁻¹ and continued for 60–120 minutes after the patient's transfer to the ICU. During this period, respiratory support was continued, and patients were observed for excessive postoperative bleeding, hemodynamic instability, tachypnea, and/or excessive respiratory muscle efforts. Sixty to 120 minutes after the end of the surgery, remifentanyl infusion was discontinued. The ventilator mode was changed to continuous positive airway pressure (set to 5 cm H₂O [0.5 kPa] and fraction of inspired oxygen 0.25). The accepted rate of breaths was more than 10 and less than 20 per minute, and the accepted tidal volume was more than 6 mL kg⁻¹. After 30 minutes of spontaneous breathing, an arterial blood gas sample was collected. The accepted pO₂ was at least 60 mm Hg, and the accepted CO₂ was less than 50 mm Hg. If all elements were fulfilled, the patient's trachea was extubated. Patient-controlled analgesia (PCA) was started via a pump that supplied incremental bolus doses of oxycodone (1 mg per bolus dose, lock-out time at seven-minute intervals, no basal infusion) during the first postoperative day. Patients were ambulated within the first 24 postoperative hours and transferred to the surgery ward by the end of the first postoperative day. The chest drain was removed approximately 10–12 hours after the patient's extubation, according to the surgeon's discretion.

In the standard care group, none of the regional anesthesia techniques were done before

TABLE 1. Patient demographics

Parameters	Standard care	ERAS	P-value
Number of patients	28	29	–
Age (years)	66.79 (63.47–70.11)	64.28 (61.67–66.88)	0.25
Body mass (kg)	80.14 (75.18–85.10)	79.83 (73.91–85.75)	0.31
Height (cm)	169.8 (166.9–172.7)	170.9 (168.0–173.9)	0.85
ASA	3 (3–3)	3 (3–3)	0.83
CCS	3 (3–3)	3 (2–3)	0.11
NYHA	1 (1–3)	1 (1–3)	0.99

The table presents patient demographics and preoperative results of ASA, CCS, and NYHA scoring systems. Patient age, body mass, and height are presented as means and confidence intervals. Statistics for these parameters were calculated with Student’s *t*-test. Results of ASA, CCS, and NYHA were presented as medians and interquartile ranges and calculated with the Mann-Whitney *U* test

ASA – American Society of Anesthesiologists, CCS – Canadian Cardiovascular Society, ERAS – Enhanced Recovery After Surgery, NYHA – New York Heart Association

the surgery. Induction of general anesthesia was performed in a manner similar to the ERAS group; however, fentanyl (1–3 µg kg⁻¹; Fentanyl WZF, Polfa Warszawa, Poland) was used instead of remifentanyl. Additional doses of fentanyl were administered according to the anesthesiologist’s discretion. Patients in the standard care group were transferred to the ICU and sedated with continuous infusion of propofol and fentanyl until the other day. Before discontinuation of mechanical ventilation, 0.1 mg kg⁻¹ of morphine (Morphini Sulfas WZF, Polfa Warszawa, Poland) was given and the PCA pump was started (1 mg per bolus dose, lock-out time at seven-minute intervals, no basal infusion).

In both groups, pain management also included one gram of intravenous paracetamol every six hours. The nurses used the numerical rating scale (NRS) at regular intervals to evaluate postoperative pain.

Outcomes

The primary outcome of this study was the postoperative mechanical ventilation time in both groups of patients. The secondary outcomes included ICU and hospital stay, postoperative drainage time, postoperative troponin T level, pain severity evaluated via the NRS, the total consumption of opioids, and perioperative complications in both groups of patients.

Pain severity was measured five times. It was first measured after the extubation, when the PCA pump was started, and then it was measured in regular six-hour intervals. The pain was measured on the NRS (0–10) by nurses who were not involved in the study.

Two different opioids, oxycodone and morphine, were used in the postoperative period; thus direct comparison of these analgesics was not possible. However, because 1 mg of morphine is comparable to 1 mg of oxycodone and the same PCA pump lock-out was set in both groups, we decided

to compare the positive demands of each patient (i.e., the number of boluses of each opioid received by patients in the standard care group and the ERAS group in 24 hours).

The current trial was planned as a feasibility study. The data obtained here will help determine the sample size for further studies.

Statistical analysis

Student’s *t*-test was used to analyze parametric data. These data were presented as means and 95% confidence intervals. Statistics derived from non-parametric data were calculated using the Mann-Whitney *U* test and are presented as medians and interquartile ranges. All measurements were performed using Statistica 13.1 software (Stat Soft, Inc., Tulsa, United States).

RESULTS

The study was conducted between November 2018 and July 2019. Overall, 57 patients were analyzed, 29 in the ERAS group and 28 in the standard care group. Only one patient declined to participate in the study. Patient demographics are presented in Table 1. No difference was found between the ERAS and the standard care groups.

Anesthesia and surgery time did not differ between the ERAS and the standard care groups. Anesthesia and surgery time was 4 (3.0–4.4) h and 3 (2.2–3.5) h in the standard care group and 4 (3.0–4.0) h and 3 (2.1–3.1) h in the ERAS group, respectively.

Primary outcome

The mechanical ventilation time was significantly shorter in the ERAS group (1 [1–3] h) than in the standard care group (10.5 [8–13.3] h; *P* = 0.00001).

Secondary outcomes

Patients in the ERAS group spent less time in the ICU (20 [16–24] h) compared to the standard care group (48 [36–48] h; *P* = 0.00001). Moreover, the ERAS patients were discharged from the hospital before the standard care patients. The mean hospital stay was 7 [6–8] days in the ERAS group and 10 [8–12] days (*P* = 0.0004) in the standard care group.

The pain intensity was significantly lower at each evaluation. The detailed pain severity description is presented in Table 2. Moreover, the PCA demand was lower in the ERAS group than in the standard care group. Patients in the ERAS group and the standard care group received 3 (3–6) and 25 (20–25) opioid boluses, *P* = 0.000001, respectively.

The postoperative drainage time was significantly shorter in the ERAS group. Thoracic drains were removed after 12 (11–17) h in the ERAS group and 23 (19–27) h (*P* = 0.00001) in the standard care

group. Postoperative drainage volume was comparable between the groups and was 480 (400–550) mL in the standard care group and 400 (300–520) mL in the ERAS group.

Postoperative troponin T concentration was significantly lower in the ERAS group (135 [94–197] ng L⁻¹) than in the standard care group (305 [240–389] ng L⁻¹; $P = 0.000061$). No difference was found between groups in the preoperative troponin level (Figure 1).

Perioperative complications

Several serious adverse events occurred in our patients, all in the standard care group. Two patients required reoperation due to severe bleeding, two had postoperative pneumonia, likely ventilatory-associated, and three developed symptoms of hyperactive delirium. No difference was noted between groups regarding the number of transfused blood products.

DISCUSSION

The results presented in the study showed the beneficial effects of ERAS protocol implementation in patients undergoing OP-CABG surgery. The time of mechanical ventilation, thoracic drainage, and ICU and hospital stay were shorter in the ERAS group than in the standard care group. The pain was less intense in the ERAS patients, and the postoperative opioid demand was reduced. Moreover, the increase of the troponin T concentration in plasma was significantly smaller in the ERAS group than in the standard care group. This suggests less myocardial damage due to ischemia-reperfusion, which inevitably occurs during the procedure. Finally, more serious complications occurred in the standard care group in the postoperative period.

We believe that these significant differences in patient outcomes were possible due to ERAS protocol implementation. Among the many elements contained in the protocol, the ESP block and remifentanyl were the most important. The feasibility of this regional blockade was illustrated in our previous study [9].

To our knowledge, the current study can be compared with two others regarding ESP block utilization in cardiac surgical patients. In the study by Krishna *et al.*, the ESP block was compared prospectively to a group without any regional analgesia technique [10]. The authors of this paper presented some beneficial effects of the ESP block, including shorter mechanical ventilation time and lower pain intensity – similar to our study's findings. However, in contrast to the current study, fentanyl was used in both groups as intraoperative and rescue analgesia in the postoperative period. Moreover, PCA with morphine or oxycodone was not a part of the study protocol.

TABLE 2. Pain severity

Timepoint	Standard care	ERAS	P-value
NRS 0	5 (4.00–5.75)	3 (2.0–4.5)	0.0001
NRS 6	4 (3.00–5.00)	2.5 (2.0–4.0)	0.004
NRS 12	4 (3.00–4.25)	3 (2.0–3.5)	0.01
NRS 18	3 (2.00–4.00)	2 (0–3.0)	0.001
NRS 24	3 (1.50–4.00)	0 (0–1.5)	0.000002

The table presents pain intensity evaluated using the numerical rating scale (NRS) where 0 = no pain, 10 = worst imaginable pain. The first assessment was done immediately after the patients' extubation when the PCA was started. The NRS was shown as medians and interquartile ranges. Probability was calculated with the Mann-Whitney *U* test. ERAS – Enhanced Recovery After Surgery, NRS – numerical rating scale




FIGURE 1. Postoperative troponin T values in the standard care and the ERAS (Enhanced Recovery After Surgery) group

Continuous epidural analgesia was compared with a continuous bilateral ESP block in the study by Nagaraja *et al.* [11]. In this article, patients in the ESP group had lower pain intensity at 24, 36, and 48 hours at rest and during coughing. Like the previously mentioned study, fentanyl was used perioperatively in both groups and PCA was not implemented in the postoperative period.

Importantly, both studies showed that the ESP block could be performed easily and did not cause any relevant complications. However, more studies are necessary to establish the role of the ESP block in cardiac surgery and compare it to other regional anesthesia techniques.

Although fentanyl is still the opioid of choice for most cardiac anesthetists, the results of numerous trials showed a positive impact of remifentanyl on postoperative troponin level and perioperative complications in cardiac surgery [12]. It seems that continuous remifentanyl infusion alleviates sympathetic input, which is caused by surgical stress [13]. Thus, fluctuations in heart rate and blood pressure are reduced in patients on remifentanyl during the surgery.

The present data confirm our previous results, suggesting possible cardioprotection provided by the combined regional/remifentanil anesthesia [14].

It appears that most of the relevant postoperative problems in the standard care group might be entailed by prolonged mechanical ventilation and sedation. Many studies have shown that both delirium and pneumonia are associated with longer mechanical ventilation under anesthesia/sedation [15, 16]. It might be reasonable to shorten sedation and mechanical ventilation time to reduce some of the perioperative complications.

The one-year experience after implementing the ERAS protocol in cardiac surgery is described in the study by Williams *et al.* [17]. The authors of this article presented a significant improvement in patient outcome after inclusion of several elements, including preoperative patient education, carbohydrate loading before general anesthesia, opioid-sparing analgesia, goal-directed perioperative insulin infusion, and bowel regimen. Most perioperative process elements that should be considered by multi-specialist cardiac surgery teams are summarized in the guidelines by Engelman *et al.* and in the recommendations by Gregory *et al.* [18, 19]. Interestingly, in both articles, regional anesthesia techniques are mentioned only as a potential part of a multimodal pain management strategy, not as routinely implemented methods.

Our study has some limitations. First, it was not a randomized controlled trial; thus, patients were recruited in the series. This increased the risk of selection bias. Moreover, the standard care group was analyzed retrospectively. More than one intervention/modification differed in both groups, including the bilateral ESP block in the ERAS group, different opioids during anesthesia and surgery, and prolonged mechanical ventilation under sedation in the standard care group. Pain intensity was evaluated in the same time interval, but the first pain assessment in the standard care group occurred much later than in the ERAS group due to the prolonged mechanical ventilation period.

CONCLUSIONS

The ERAS protocol implementation in OP-CABG surgery can improve patient outcomes. This rationale is supported by the results presented in the current study, including the shorter time of mechanical ventilation, thoracic drainage, and ICU and hospital stay. Moreover, pain management was improved in patients who were treated according to the ERAS protocol. Finally, the new strategy might decrease postoperative troponins and reduce complication rates. Our study confirmed that introduction of the ERAS protocol was feasible and would allow for a fully-powered study.

ACKNOWLEDGEMENTS

1. Financial support and sponsorship: none.
2. Conflict of interest: none.

REFERENCES

1. Weiss AJ, Elixhauser A. Trends in Operating Room Procedures in U.S. Hospitals, 2001-2011: Statistical Brief #171. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD): Agency for Healthcare Research and Quality (US), 2006.
2. Head SJ, Milojevic M, Taggart DP, Puskas JD. Current practice of state-of-the-art surgical coronary revascularization. *Circulation* 2017; 136: 1331-1345. doi: 10.1161/CIRCULATIONAHA.116.022572.
3. Moazzami K, Dolmatova E, Maher J, et al. In-hospital outcomes and complications of coronary artery bypass grafting in the United States between 2008 and 2012. *J Cardiothorac Vasc Anesth* 2017; 31: 19-25. doi: 10.1053/j.jvca.2016.08.008.
4. Stephens RS, Whitman GJ. Postoperative critical care of the adult cardiac surgical patient. Part I: routine postoperative care. *Crit Care Med* 2015; 43: 1477-1497. doi: 10.1097/CCM.0000000000001059.
5. Zubrzycki M, Liebold A, Skrabal C, et al. Assessment and pathophysiology of pain in cardiac surgery. *J Pain Res* 2018; 11: 1599-1611. doi: 10.2147/JPR.S162067.
6. Caruso TJ, Lawrence K, Tsui BCH. Regional anesthesia for cardiac surgery. *Curr Opin Anaesthesiol* 2019; 32: 674-682. doi: 10.1097/ACO.0000000000000769.
7. Mietto C, Pinciroli R, Patel N, Berra L. Ventilator associated pneumonia: evolving definitions and preventive strategies. *Respir Care* 2013; 58: 990-1007. doi: 10.4187/respcare.02380.
8. Ljungqvist O, Scott M, Fearon KC. Enhanced recovery after surgery: a review. *JAMA Surg* 2017; 152: 249-252. doi: 10.1007/s40520-018-0905-1.
9. Borys M, Gawęda B, Horeczy B, et al. Erector spinae-plane block as an analgesic alternative in patients undergoing mitral and/or tricuspid valve repair through a right mini-thoracotomy – an observational cohort study. *Videosurgery Miniinv* 2020; 15: 208-214. doi: 10.5114/wiitm.2019.85396.
10. Krishna SN, Chauhan S, Bhoi D, et al. Bilateral erector spinae plane block for acute post-surgical pain in adult cardiac surgical patients: a randomized controlled trial. *J Cardiothorac Vasc Anesth* 2019; 33: 368-375. doi: 10.1053/j.jvca.2018.05.050.
11. Nagaraja PS, Ragavendran S, Singh NG, et al. Comparison of continuous thoracic epidural analgesia with bilateral erector spinae plane block for perioperative pain management in cardiac surgery. *Ann Card Anaesth* 2018; 21: 323-327. doi: 10.4103/aca.ACA_16_18.
12. Greco M, Landoni G, Biondi-Zocai G, et al. Remifentanil in cardiac surgery: a meta-analysis of randomized controlled trials. *J Cardiothorac Vasc Anesth* 2012; 26: 110-116. doi: 10.1053/j.jvca.2011.05.007.
13. Shu A, Zhan L, Fang H, et al. Evaluation of remifentanil anesthesia for off-pump coronary artery bypass grafting surgery using heart rate variability. *Exp Ther Med* 2013; 6: 253-259. doi: 10.3892/etm.2013.1108.
14. Kurowicki A, Borys M, Żurek S, et al. Remifentanil and sevoflurane based anesthesia combined with bilateral erector spinae plane block in patients undergoing off-pump coronary artery bypass graft surgery. *Videosurgery Miniinv* 2019. doi: 10.5114/wiitm.2019.88748.
15. Reade MC, Finfer S. Sedation and delirium in the intensive care unit. *N Engl J Med* 2014; 370: 444-454. doi: 10.1056/NEJMra1208705.
16. Timsit JF, Esaied W, Neuville M, Bouadma L, Mourvillier B. Update on ventilator-associated pneumonia. *F1000Res* 2017; 6: 2061. doi: 10.12688/f1000research.12222.1.
17. Williams JB, McConnell G, Allender JE, et al. One-year results from the first US-based enhanced recovery after cardiac surgery (ERAS Cardiac) program. *J Thorac Cardiovasc Surg* 2019; 157: 1881-1888. doi: 10.1016/j.jtcvs.2018.10.164.
18. Engelman DT, Ben Ali W, Williams JB, et al. Guidelines for perioperative care in cardiac surgery: enhanced recovery after surgery society recommendations. *JAMA Surg* 2019; 154: 755-766. doi: <https://doi.org/10.1001/jamasurg.2019.1153>.
19. Gregory AJ, Grant MC, Manning MW, et al. Enhanced Recovery After Cardiac Surgery (ERAS Cardiac) recommendations: an important first step-but there is much work to be done. *J Cardiothorac Vasc Anesth* 2020; 34: 39-47. doi: 10.1053/j.jvca.2019.09.002.