

Serum reactive oxygen species and apoptosis markers in septic patients

Sebastián Pablo Chapela^{1,2}, Isabel Burgos¹, Amalia Schiel³, Manuel Alonso⁴, Carlos Alberto Stella¹

¹Universidad de Buenos Aires, Facultad de Medicina, Departamento de Bioquímica Humana (INBIOMED UBA-CONICET), Argentina

²Servicio de Terapia Intensiva, Hospital Británico de Buenos Aires, Argentina

³Laboratorio Central, Hospital Británico de Buenos Aires, Argentina

⁴Universidad de Buenos Aires, Ciclo Básico Común, Departamento de Ciencias Biológicas, Argentina

Abstract

Background: Oxidative stress is one of the pathophysiological processes that occur during sepsis. Reactive oxygen species (ROS) production causes lipid peroxidation and protein and DNA damage. ROS and DNA damage triggers apoptosis. Several studies have shown that organ failure in sepsis is mediated by apoptosis. The aim of this study is to investigate the levels of serum ROS and serum caspase-3 in septic patients and healthy volunteers, and their correlation.

Methods: Serum samples were taken within the first 12 hours of ICU stay. The dichlorofluorescein technique was used to determine serum ROS levels, and the ELISA technique was used to quantify serum caspase-3 in septic patients and healthy volunteers.

Results: There was no difference in serum ROS levels between healthy volunteers and septic patients ($P = 0.26$), and there was a significant difference in serum caspase-3 levels between healthy volunteers and septic patients ($P < 0.001$). There was no difference between patients who lived and died in the intensive care unit (ICU) in serum ROS ($P = 0.089$) and serum caspase-3 ($P = 0.18$). There was no correlation between both markers ($R = -0.0013$, $P = 0.98$).

Conclusions: We conclude that there is no correlation between serum ROS and caspase-3; therefore, both processes might not be associated during the first hours of ICU stay.

Key words: reactive oxygen species, apoptosis, caspase, sepsis.

Anestezjologia Intensywna Terapia 2021;
53, 2: 126–133

Otrzymano: 31.05.2020,
zaakceptowano: 26.12.2020

ADRES DO KORESPONDENCJI:

Sebastián Pablo Chapela, Departamento de Bioquímica Humana, Facultad de Medicina, Universidad de Buenos Aires, Paraguay 2055, CABA, Argentina, ZIP 1121, e-mail: spchapela@hotmail.com

Sepsis is a syndrome of physiological, pathological, and biochemical abnormalities, with several pathophysiological processes that occur simultaneously [1]. The worldwide incidence of sepsis is increasing [2–6], and therefore the correct understanding of its pathophysiology, diagnosis, and treatment are important. Depending on the definition used, incidence in the United States varies between 300 per 100,000 and 1031 per 100,000 inhabitants, with an annual growth of 13% [4]. Hospital mortality varies between 14.7% and 29% [4]. In some studies, carried out in Latin-America, it is even higher [7–10].

Among the pathophysiological processes, mitochondrial dysfunction, which is caused by a decreased function of certain complexes in the electron transport chain, producing less energy and higher levels of free radicals, can be found. *In vivo*, reactive oxygen species (ROS) are produced in different pathways, different cells, and in different quantities. In severe pathologies, phagocytes are

the main cells that produce ROS as an element of immune defence, and their purpose is the destruction of microorganisms. Under normal circumstances, there is a continuous production of small amounts of ROS in the mitochondria and, more precisely, in the electron transport chain [11]. It is believed that the latter is responsible for 90% of ROS produced in the cell [12, 13]. Immune system and mitochondrial dysfunction are the main sources of ROS. In sepsis, ROS and reactive nitrogen species (RNS) are increased, both in circulation and in tissue [14].

The plasma oxidative burst takes place in the early stages of sepsis and is carried out by neutrophils, macrophages, and endothelial cells. Antioxidants remove ROS from the plasma, which leads to a decrease in the concentration of these molecules [15]. The level of free thiols decreases due to oxidation or nitrosylation [15]. As a result, the level of antioxidant capacity of plasma is reduced [15]. There are several proteins that activate signalling pathways that promote apoptosis and are sensitive

to superoxide [16]. Also, hydroxyl radical is a reactive species that oxidizes lipids, proteins, and DNA, which results in genomic instability [17].

On the other hand, another pathophysiological process described in sepsis is apoptosis. Apoptosis is a highly regulated and conserved cell death process during which the cell self-destructs [16]. It is a normal process, which, in multicellular organisms, eliminates undesired or superfluous cells, neutralizing the potential damage caused by cells with defective DNA [16].

Cells in which apoptosis is triggered decrease in size and are condensed and fragmented, releasing membrane-limited apoptotic bodies that are generally absorbed by other cells [18]. In addition, the nucleus is condensed, the DNA is fragmented, and the cellular constituents are not released to the extracellular medium where they could harm the surrounding cells, in contrast with what happens in cell death by necrosis [18]. Apoptosis involves enzymes that cleave proteins at the cysteine amino acid level (caspases) [18]. Most, but not all, of the apoptotic pathways lead to the activation of these enzymes [16].

Different triggers have been proposed as the cause of organ failure: Decreased tissue perfusion and its resulting respiratory failure, mitochondrial dysfunction, the production of ROS, and other mechanisms, cell death being the final outcome [19].

In recent years, interest in apoptosis as a mechanism of cell death has increased. There are studies that show increased levels of apoptosis at the cellular level in different models of sepsis [20–23], but there is insufficient evidence regarding systemic apoptosis levels in septic patients, or the relationship with mortality in intensive care units (ICU) [24, 25]. While several organs show evidence of inflammation with migration of inflammatory cells, increased interstitial fluid, increased capillary permeability, and epithelial disruption, the degree of cell death is low [26]. The degree of cell death is disproportionately low compared with the degree of clinical severity or the biochemical presentation of organ dysfunction [26, 27]. Also, there is indirect evidence of lymphocyte apoptosis in septic patients with extensive depletion of lymphocytes in the white pulp of the spleen [28].

Both pathophysiological processes (ROS production and apoptosis) and others like inflammation, thrombosis, microabscess formation, and ischaemic changes are causes of the multiple organ dysfunction observed in sepsis, which, if left uncontrolled, leads to patient death [29]. The aim of this study was to investigate the level of serum markers of ROS and apoptosis and to ascertain whether those processes occur simultaneously. There is no literature relating ROS levels and apoptosis in sepsis.

METHODS

Study design

This is a prospective observational trial, carried out in a single centre. The centre is a private tertiary hospital with an intensive care unit with 27 beds, in Buenos Aires, Argentina. Subjects were included between May 2013 and December 2017. The protocol was approved by the local internal review board (number of approval: CRIHB#521). All patients signed an informed consent form.

Inclusion and exclusion criteria

Inclusion criteria for the patients were as follows: aged between 18 and 80 years with a diagnosis of sepsis, according to the Sepsis-2 definition [30]. Inclusion criteria for the healthy volunteers were as follows: over 18 years of age with no actual septic pathology or signs. Exclusion criteria for the septic patients were the following: refusal of informed consent, limitation of therapeutic efforts at the time of inclusion in the study, and pregnancy.

Treatment of sepsis

The diagnosis and treatment of sepsis was decided by ICU physicians. For treatment, the recommendations of the 2013 and 2016 “Surviving Sepsis Campaign” guides were followed [2, 31].

Measurement of serum reactive oxygen species

Upon arrival of the patient, within 12 hours of admission to the ICU, venous blood samples were taken. After that, blood was centrifuged for 5 minutes at 3000 rpm, and serum was frozen at -70°C , as described in the bibliography [32–34]. Using dichlorofluorescein-diacetate (DCFH) (Sigma Aldrich. Catalogue D6665) ROS were measured. 12 μL of serum were incubated for 10 minutes in 1000 μL of TE buffer, and 10 μL of NaOH and 10 μL of DCFH solution were added. Fluorescence was measured after creating an emission spectrum between 500 and 550 nm with each serum sample; the peak of emission at 525 nm was recorded.

Measurement of serum levels of caspase-3

In the same sample described for the measurement of serum ROS, the quantification of caspase-3 levels was carried out using the ELISA method on patients' serum. The kit was manufactured by MyBioSource[®] – Human Caspase-3 ELISA KIT Catalogue #MBS040290, batches #02/2016 and #10/2017. In order to do this, the patients' serum, frozen at -70°C , was used. The levels of caspase-3 are shown in pg mL^{-1} . Samples were measured using an EMP M201[®] Microplate Reader (China).

Statistical analysis

Normal and non-normal distribution data were determined based on mean, median, and kurtosis. The mean was calculated with standard deviation (SD) to describe quantitative variables in normal distribution data, and with 95% confidence interval (95% CI) in non-normal data. In order to analyse whether there were significant differences in continuous variables between two groups, Student's *t*-test was used for variables with normal distribution, and the *U* Mann-Witney test was used for variables with abnormal distribution. To compare nominal and ordinal qualitative variables, the chi-square test was used. To correlate two variables, the Spearman test was used, depending on whether the distribution of the variables was normal or not. In all cases, a significance level of $P < 0.05$ was considered. The following statistical programs were used: EPIinfo 7.0, Statistix 7.0 (Informer Technologies, Inc., Atlanta, USA), and Graph Pad Prism 8.0.2. (Graph-Pad Software, San Diego, USA).

RESULTS

Table 1 shows the comparison of demographic and clinical parameters between patients and healthy volunteers. A total of 124 people were enrolled, 98 of whom were patients with severe sepsis or septic shock diagnoses, and 26 were healthy volunteers. None of the healthy volunteers had a chronic disease that required treatment. The healthy volunteers were younger than the patients. Table 2 describes the severity of disease scores and origin of infection.

Serum ROS were measured using the DCFH technique. There were no differences in the ROS levels between healthy volunteers and septic patients (Emission = 0.1077 [SD = 0.0324] vs. 0.108 [SD = 0.0364]; $P = 0.26$) (Figure 1A). Twenty-eight out of the 98 patients died during hospitalization. When analysing only septic patients, there were no differences in the ROS levels of patients who survived and those who died in the ICU (Emission = 0.1103 [SD = 0.0384] vs. 0.1022 [SD = 0.0303]; $P = 0.089$) (Figure 1B). Finally, 7 patients died during the first day of hospitalization, and there were no differences between those patients and the patients who survived (Emission = 0.1076 [SD = 0.0369] vs. 0.1145 [0.029]; $P = 0.31$) (Figure 1C).

Serum levels of caspase-3 were higher than those of healthy volunteers (31.75 pg mL⁻¹ [95% CI: 25.84–37.65] vs. 7.67 pg mL⁻¹ [95% CI: 1.93–13.4]; $P < 0.001$) (Figure 2A). When analysing only patients, there were no statistically significant differences between patients who lived and those who died during their hospitalization in the ICU (28.49 pg mL⁻¹ [95% CI: 22.21–34.77] vs. 39.89 pg mL⁻¹

TABLE 1. Description of the population enrolled in this study

Parameter	Healthy volunteers, <i>n</i> = 26	Patients, <i>n</i> = 98	<i>P</i> -value
Age, years	49	64.38	< 0.001
Sex (male/female), <i>n</i>	11/15	56/42	0.25
Tobacco use, <i>n</i>	2	56	< 0.001
Diabetes, <i>n</i>	2	54	< 0.001
Hypertension, <i>n</i>	2	48	< 0.001
Dyslipidaemia, <i>n</i>	3	18	0.56
Ischemic cardiopathy, <i>n</i>	0	15	0.038
Heart failure, <i>n</i>	0	6	0.34
COPD, <i>n</i>	0	9	0.20
Atrial fibrillation, <i>n</i>	0	9	0.20
Chronic kidney disease, <i>n</i>	0	13	0.06
Rheumatic disorder, <i>n</i>	0	8	0.20
Oncologic disease, <i>n</i>	0	24	0.002
Oncohaematological disease, <i>n</i>	0	11	0.45
HIV, <i>n</i>	0	2	1

The χ^2 and Student tests were used to determine whether there were differences between the groups.

[95% CI: 26.22–53.57]; $P = 0.18$) (Figure 2B). Finally, there were no significant differences in the levels of caspase-3 between patients who survived and those who died on the first day (29.83 pg mL⁻¹ [95% CI: 23.99–35.67] vs. 56.69 pg mL⁻¹ [95% CI: 21.82–91.56]; $P = 0.1$) (Figure 2C).

The Spearman test was performed to demonstrate any correlation between DCFH emission levels and serum levels of caspase-3 of septic patients and those of healthy volunteers. There was no correlation between the levels of both variables. A scatter plot with levels of both variables was made (Figure 3). In addition, the Spearman test was also performed to demonstrate any correlation between DCFH emission levels and serum levels of caspase-3

TABLE 2. Description of the septic patient population. The characteristics of the septic patient population are described. The APACHE II score values were taken with the worst result in the first 24 hours and the SOFA score with the values at admission

Origin of infection	<i>n</i> (%)
Abdominal	34 (35.5)
Respiratory	26 (26.28)
Urinary	15 (15.46)
Endovascular	6 (6.19)
Skin and muscle	9 (9.28)
Neurological	2 (2.06)
Other	6 (6.19)
APACHE II	17.35 (SD: 7.26)
SOFA	7.33 (SD: 3.11)




FIGURE 1. Serum reactive oxygen species (ROS) levels. **A)** ROS levels in healthy volunteers and patients. Dichlorofluorescein-diacetate (DCFH) emission levels are stated at 525 nm as a marker of ROS. There was no significant difference between groups. Student's test ($P = 0.26$). **B)** ROS levels of patients who lived and died in the ICU. DCFH emission levels are stated at 525 nm as a marker. There was no significant difference between groups. Student's test ($P = 0.089$). **C)** ROS levels in septic patients who lived and those who died on the first day of hospitalization. DCFH emission levels are stated at 525 nm as a marker of ROS. There was no significant difference between groups. Student's test ($P = 0.31$)




FIGURE 2. Serum apoptosis levels. **A)** Serum levels of Caspase-3 in healthy volunteers and septic patients. There were significant differences between groups when using the Mann Whitney test ($P < 0.001$). **B)** Serum levels of Caspase-3 in patients who lived and those who died in the ICU. There were no significant differences between groups. Mann Whitney test ($P = 0.18$). **C)** Serum levels of Caspase-3 in patients who lived and those who died during the first day of hospitalization. There were no significant differences between groups. Mann Whitney test ($P = 0.1$)




FIGURE 3. Correlation between reactive oxygen species (ROS) levels and serum levels of caspase-3. ROS were measured using the dichlorofluorescein-diacetate (DCFH) technique and serum levels of caspase-3 were measured in healthy volunteers and controls. There was no correlation between both variables ($R = -0.0013$, $P = 0.98$)

only among septic patients. There was no correlation between the levels of both variables. A scatter plot with levels of both variables was constructed (Figure 4).

Finally, correlation between serum ROS and serum caspase-3 levels with the severity scores SOFA and APACHE II was also analysed. There was no correlation between the levels of caspase-3 and the SOFA score ($R = 0.1823$; $P = 0.08$), nor with the APACHE II score ($R = 0.1047$; $P = 0.32$). There was also no correlation between APACHE II levels and ROS

levels ($R = -0.0562$; $P = 0.6$). In contrast, there was an inverse correlation between SOFA levels and activated ROS levels ($R = -0.2146$; $P = 0.04$).

DISCUSSION

In this study, we included 98 septic patients and 26 healthy volunteers. All patients were admitted to the ICU with severe sepsis and septic shock diagnoses, in accordance with the second definition of sepsis, which was valid at the start of the study [30]. However, considering the Sepsis-3 consensus [1], the same group of patients would have been included. This study is innovative because a group of healthy volunteers were included, and serum ROS and caspase-3 levels were obtained from them.

It was noted that there were no differences in systemic ROS levels between healthy volunteers and septic patients, and no differences between septic patients who died ICU and those who survived. Moreover, there were no differences between patients who died during the first day of hospitalization and those who survived. These results are not in accordance with those obtained in previous studies performed on rats, in which the septic group had higher ROS levels than the control group, using the same technique to measure ROS [35]. These differences may be due to methodological issues because, at the time of sample collection, the rats had

had sepsis for 24 h, while patients in the hospital came in through the emergency service, they were supported with fluids, the first dose of antibiotics was administered, then they were admitted to the ICU, a central line was placed, they were intubated if necessary, vasoactive drugs were administered, and finally the samples were taken. Another possible cause for this difference between the results obtained from rats and those from patients is that they are different models, with a different pathophysiology, and therefore it is often difficult to replicate animal results in patients [15, 29].

This study is innovative because serum ROS were measured in septic patients instead of oxidative stress markers. Previous literature associated damage at the tissue level with ROS [36–41]. Using the current technique, ROS levels in yeast were measured under stress conditions with Menadione [42]. The DCFH technique is an already described technique to measure intracellular ROS also in extracellular fluids [43–46]. It has been demonstrated in our laboratory that septic rats have higher levels of ROS than controls, and that the administration of parental succinate decreases ROS levels [35].

In several models, it has been shown that ROS damage cell structures including DNA, which leads to apoptosis. The presence of serum ROS in an animal model of sepsis are also increased, but this is not associated with markers of organ failure [47]. Lorente *et al.* [48] showed that the total antioxidant capacity of septic patients who did not survive was higher and, in turn, this was a marker of 30-day mortality. Also, septic patients with a SOFA score higher than 7 had higher ROS levels measured with DCFH [49].

DCFH emission results and corresponding serum ROS levels in septic patients are not in accordance with studies that measure oxidative stress, probably because different techniques are used. In this study, DCFH emission that states serum ROS levels was measured, while in other studies, either antioxidant capacity or oxidative stress substitutes, such as TBARS or oxidized proteins, were measured.

On the other hand, apart from the measurement of ROS and oxidative stress markers, multiple studies with antioxidants have been carried out. Preclinical and clinical trials with antioxidants had diverse results. In a preclinical murine model of sepsis, circulating cytokines, increased urea, ALT, and LDH, decreased glucose, lymphopaenia/neutrophilia, and NO blood release were not altered with the treatment with SkQ1 antioxidants, and there was no difference in survival with antioxidant mtAOX [50]. Other preclinical trials had positive results. Superoxide dismutase (SOD), catalase (CAT), and ascorbate peroxidase (APX) activities were decreased after the administration of antioxidant quercetin in septic




FIGURE 4. Correlation between systemic reactive oxygen species (ROS) levels and serum levels of caspase-3 in septic patients. There was no correlation between both variables ($R = -0.0732$, $P = 0.47$)

rats [51]. In rats that were injected intraperitoneally with lipopolysaccharide, it was observed that pre-treatment with N-acetylcysteine (NAC) decreased certain parameters of oxidative stress [52, 53]. Also, in a study with septic rats, the administration of intraperitoneal NAC, associated with deferoxamine, decreased the production of superoxide, the activity of enzyme catalase, superoxide dismutase, and myeloperoxidase in all organs, in addition to improving survival [54]. In several clinical trials in which septic patients were supplemented with antioxidants, the length of mechanical ventilation or oxygen levels were not reduced [55–68]. There are many works that describe differences between animal models of sepsis and patients, and the difficulty of transferring the results of the animal model to patients [69–71]. The absence of positive results in clinical studies with antioxidants has some sort of relationship with the results of the current study, in which there were no differences between the ROS levels of healthy volunteers and septic patients.

In addition, septic patients had higher serum levels of caspase-3 than healthy volunteers, with statistically significant differences. The median serum levels of caspase-3 in patients who died was 75% higher than in those who survived, but the differences were not statistically significant. Similarly, the median of patients who died on the first day was 90% higher than those who survived the first day, but these differences were not statistically significant. This is due to the statistical test chosen and the spread of data with a large confidence interval width [72], but it does not make the data less relevant [73].

When comparing the data obtained with those from the literature, there are certain similarities. In a study carried out by Lorente *et al.* [74] a higher mortality in septic patients with higher serum levels of caspase-3 measured using the ELISA technique was shown. In this study, higher caspase-3 levels

are associated with higher mortality (odds ratio, 6.51; 95% CI: 3.32–12.77; $P < 0.001$) [74]. In the same study, a positive correlation between caspase-3 and a different marker of apoptosis, caspase-cleaved cytokeratin-18, was found [74]. The same group published a study in which caspase-cleaved cytokeratin-18 was higher in septic patients who died in the ICU and had a positive correlation with levels of interleukin 6, APACHE II, SOFA, and lactate [75]. Similarly, this research group found that higher levels of caspase-cleaved cytokeratin-18 (levels over 20 UI L^{-1}) resulted in higher mortality (OR = 8.476; 95% CI: 2.087–34.434; $P = 0.003$) [76]. Although these results contrast with those obtained in the present work, it should be taken into account that patients were different, and the laboratory tests to measure serum caspase were different. These are possible explanations for these differences in the results.

Also, this study is innovative because it was noted that there was no correlation between serum levels of ROS and caspase-3 between healthy volunteers and septic patients. Nor was there a correlation between serum levels of ROS and caspase-3 when analysing only the patient group. This is an interesting finding, which suggests that there is no association between those molecular processes in sepsis. No other study in the bibliography that quantified ROS and apoptosis markers was found. ROS are molecules with a very short life. A limitation of this study is that samples were taken in the ICU, after a central line was placed and vasoactive drugs were indicated. More studies with samples taken at hospital admission and different times of stay should be performed to clarify this finding.

Finally, it should be clarified first that, although the recruitment time was relatively long, the conditions remained the same during the study and all patients were treated by the same medical team, although the therapies could have changed slightly in that period. Second, only ROS were measured in serum; other measurements could have been carried out (i.e. carbonylated proteins or TBARS), but the objective was to observe the effect of the relationship between these chemical species and mortality, as well as their relationship with apoptosis markers. Finally, it should be considered that the severity of the patients' disease was very heterogeneous, because obtaining the largest possible number of cases was deemed more important than the homogeneity of the sample. A more homogeneous sample could have generated other results.

CONCLUSIONS

In the present work, serum ROS levels were measured, and septic patients did not show higher se-

rum ROS levels than healthy volunteers. Conversely, patients showed higher serum levels of caspase-3 than healthy volunteers. There was no correlation between the serum levels of both markers. These results would indicate that serum ROS do not serve as markers of oxidative stress in septic patients, despite the fact that ROS production is known to take place at the tissue level, but instead, serum apoptosis markers could reveal the occurrence of cellular damage and consequent multi-organ failure in these patients relatively fast and with ease.

ACKNOWLEDGEMENTS

1. Financial support and sponsorship: This work was supported by the Universidad de Buenos Aires with the grant RS (CS) N° 8456/2017.
2. Conflicts of interest: none.

REFERENCES

1. Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016; 315: 801-810. doi: 10.1001/jama.2016.0287.
2. Dellinger RP, Levy MM, Rhodes A, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med* 2013; 39: 165-228. doi: 10.1007/s00134-012-2769-8.
3. Shankar-Hari M, Phillips G, Levy ML, et al. Developing a new definition and assessing new clinical criteria for septic shock for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016; 315: 775-787. doi: 10.1001/jama.2016.0289.
4. Gaieski DF, Edwards JM, Kallan MJ, Carr BG. Benchmarking the Incidence and Mortality of Severe Sepsis in the United States. *Crit Care Med* 2013; 41: 1167-1174. doi: 10.1097/CCM.0b013e31827c09f8.
5. Vincent J, Marshall JC, Namendys-silva SA, et al. Assessment of the worldwide burden of critical illness: the Intensive Care Over Nations (ICON) audit. *Lancet Respir Med* 2014; 2: 380-386. doi: 10.1016/S2213-2600(14)70061-X.
6. Fleischmann C, Scherag A, Adhikari NKJ, et al. Assessment of global incidence and mortality of hospital-treated sepsis – current estimates and limitations. *Am J Respir Crit Care Med* 2015; 193: 259-272.
7. Machado FR, Cavalcanti AB, Bozza FA, et al. The epidemiology of sepsis in Brazilian intensive care units (the Sepsis PREvalence Assessment Database, SPREAD): an observational study. *Lancet Infect Dis* 2017; 17: 1180-1189. doi: 10.1016/S1473-3099(17)30322-5.
8. Schultz MJ, Dunser MW, Dondorp AM, et al. Current challenges in the management of sepsis in ICUs in resource-poor settings and suggestions for the future. *Intensive Care Med* 2017; 43: 612-624. doi: 10.1007/s00134-017-4750-z.
9. Fleischmann C, Scherag A, Adhikari NKJ, et al. Assessment of global incidence and mortality of hospital-treated sepsis – current estimates and limitations. *Am J Respir Crit Care Med* 2015; 193: 259-272. The same as [6].
10. Estenssoro E, Kanoore Edul VS, Loudet CI, et al. Predictive Validity of Sepsis-3 Definitions and Sepsis Outcomes in Critically Ill Patients: A Cohort Study in 49 ICUs in Argentina. *Crit Care Med* 2018; 46: 1276-1283. doi: 10.1097/CCM.0000000000003208.
11. Preiser JC. Oxidative stress. *JPEN J Parenter Enteral Nutr* 2012; 36: 147-154. doi: 10.1177/0148607111434963.
12. Bouchez C, Devin A. Mitochondrial biogenesis and mitochondrial reactive oxygen species (ROS): a complex relationship regulated by the cAMP/PKA signaling pathway. *Cells* 2019; 8: 287. doi: 10.3390/cells8040287.
13. Russell EG, Cotter TG. New insight into the role of reactive oxygen species (ROS) in cellular signal-transduction processes. *Int Rev Cell Mol Biol* 2015; 319: 221-254. doi: 10.1016/bs.ircmb.2015.07.004.
14. Heyland DK, Dhaliwal R, Suchner U, Berger MM. Antioxidant nutrients: a systematic review of trace elements and vitamins in the critically ill patient. *Intensive Care Med* 2005; 31: 327-337. doi: 10.1007/s00134-004-2522-z.

15. Andrades M, Morina A, Spasić S, Spasojević I. Bench-to-bedside review: sepsis – from the redox point of view. *Crit Care* 2011; 15: 230. doi: 10.1186/cc10334.
16. Redza-Dutordoir M, Averill-bates DA. Activation of apoptosis signaling pathways by reactive oxygen species. *Biochim Biophys Acta* 2016; 1863: 2977-2992. doi: 10.1016/j.bbamcr.2016.09.012.
17. Fomby P, Cherlin AJ. ROS function in redox signaling and oxidative stress. *Curr Biol* 2014; 24: R453-462. doi: 10.1016/j.cub.2014.03.034.
18. Lodish H, Berk A, Matsudaira P, et al. *Biología celular y molecular. Panamericana*, 2013.
19. Aziz M, Jacob A, Wang P. Revisiting caspases in sepsis. *Cell Death Dis* 2014; 5: e1526. doi: 10.1038/cddis.2014.488.
20. Bayir H, Kagan VE. Bench-to-bedside review: mitochondrial injury, oxidative stress and apoptosis – there is nothing more practical than a good theory. *Crit Care* 2008; 12: 206. doi: 10.1186/cc6779.
21. Conus S, Perozzo R, Reinheckel T, et al. Caspase-8 is activated by cathepsin D initiating neutrophil apoptosis during the resolution of inflammation. *J Exp Med* 2008; 205: 685-698. doi: 10.1084/jem.20072152.
22. Comim CM, Barichello T, Grandgirard D, Dal-Pizzol F, Quevedo J, Leib SL. Caspase-3 mediates in part hippocampal apoptosis in sepsis. *Mol Neurobiol* 2013; 47: 394-398.
23. Circu ML, Aw TY. Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med* 2010; 48: 749-762. doi: 10.1016/j.freeradbiomed.2009.12.022.
24. Chung CS, Venet F, Chen Y, et al. Deficiency of bid protein reduces sepsis-induced apoptosis, inflammation and improves septic survival. *Shock* 2010; 34: 150-161. doi: 10.1097/SHK.0b013e3181cf70fb.
25. Exline MC, Justiniano S, Hollyfield JL, et al. Microvesicular caspase-1 mediates lymphocyte apoptosis in sepsis. *PLoS One* 2014; 9: e90968. doi: 10.1371/journal.pone.0090968.
26. Delogu G, Famularo G, Tellan G, et al. Lymphocyte apoptosis, caspase activation and inflammatory response in septic shock. *Infection* 2008; 36: 485-487.
27. Singer M. The role of mitochondrial dysfunction in sepsis-induced multi-organ failure. *Virulence* 2014; 5: 66-72.
28. Hotchkiss R, Swanson P, Freeman B, Tinsley K, Cobb J, Matuschak G, et al. Apoptotic cell death in patients with sepsis, shock, and multiple organ dysfunction. *Crit Care Med* 1999; 27: 1230-1251. doi: 10.4161/viru.26907.
29. Garofalo AM, Lorente-Ros M, Goncalvez G, et al. Histopathological changes of organ dysfunction in sepsis. *Intensive Care Med* 2019; 7 (Suppl 1): 45. doi: 10.1186/s40635-019-0236-3.
30. Levy MM, Fink MP, Marshall JC, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med* 2003; 31: 1250-1256. doi: 10.1097/01.CCM.0000050454.01978.3B.
31. Rhodes A, Evans LE, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med* 2016; 43: 304-377.
32. Lhommeau I, Douillard S, Foursac A, Aillet L, Bigot E, Patrice T. Influence of Animal Species on Secondary Oxidant in Serum exposed to Singlet Oxygen. *ILAR J* 2012; 53: e31-e42. doi: 10.1093/ilar.53.1.31.
33. Hayashi I, Morishita Y, Imai K, Nakamura M, Nakachi K, Hayashi T. High-throughput spectrophotometric assay of reactive oxygen species in serum. *Mutat Res* 2007; 631: 55-61. doi: 10.1016/j.mrgentox.2007.04.006.
34. Douillard S, Lhommeau I, Foursac A, Aillet L, Bigot E, Patrice T. Biophysical parameters influencing secondary oxidants activation in human serum exposed to singlet oxygen. *J Photochem Photobiol B Biol* 2011; 102: 224-231. doi: 10.1016/j.jphotobiol.2010.12.007.
35. Chapela SP, Burgos I, Congost C, et al. Parenteral succinate reduces systemic ROS production in septic rats, but it does not reduce creatinine levels. *Oxid Med Cell Longev* 2018; 2018: 1928945.
36. Leff JA, Parsons PE, Day CE, et al. Serum antioxidants as predictors of adult respiratory distress syndrome in patients with sepsis. *Lancet* 1993; 341: 777-780.
37. Cowley HC, Bacon PJ, Goode HF, Webster NR, Jones JG, Menon DK. Plasma antioxidant potential in severe sepsis: a comparison of survivors and nonsurvivors. *Crit Care Med* 1996; 24: 1179-1183. doi: 10.1097/00003246-199607000-00019.
38. Pavlakou P, Liakopoulos V, Eleftheriadis T, Mitsis M, Dounousi E. Oxidative stress and acute kidney injury in critical illness: pathophysiologic mechanisms – biomarkers – interventions, and future perspectives. *Oxid Med Cell Longev* 2017; 2017: 6193694.
39. Yamaguchi J, Nagase M, Yamamoto Y, et al. Increased oxidative stress and renal injury in patients with sepsis. *J Clin Biochem Nutr* 2018; 63: 137-143. doi: 10.3164/jcbn.17-130.
40. Sandesc M, Rogobete AF, Bedreag OH, et al. Analysis of oxidative stress-related markers in critically ill polytrauma patients: an observational prospective single-center study. *Bosnial J* 2017; 18: 191-197.
41. Gebicki JM. Oxidative stress, free radicals and protein peroxides. *Arch Biochem Biophys* 2016; 595: 33-39. doi: 10.1016/j.ab.2015.10.021.
42. Gamondi O, Chapela S, Nieves I, Burgos I, Alonso M, Stella C. Idenbenone treatment mediates the effect of menadione oxidative stress damage in *Saccharomyces cerevisiae*. *Drug Metab Lett* 2012; 6: 120-123.
43. Tournebize J, Sapin-minet A, Bartosz G, Leroy P. Pitfalls of assays devoted to evaluation of oxidative stress induced by inorganic nanoparticles. *Talanta* 2013; 116: 753-763. doi: 10.1016/j.talanta.2013.07.077.
44. Kuznetsov AV, Kehrler I, Kozlov AV, et al. Mitochondrial ROS production under cellular stress: Comparison of different detection methods. *Anal Bioanal Chem* 2011; 400: 2383-2390. doi: 10.1007/s00216-011-4764-2.
45. Tominaga H, Katoh H, Odagiri K, et al. Different effects of palmitoyl-L-carnitine and palmitoyl-CoA on mitochondrial function in rat ventricular myocytes. *Am J Physiol Hear Circ Physiol* 2008; 295: H105-112.
46. Tetz LM, Kamau PW, Cheng AA, Meeker JD, Loch-Carus R. Troubleshooting the dichlorofluorescein assay to avoid in measurement of toxicant-stimulated cellular production of reactive oxygen species. *J Pharmacol Toxicol Methods* 2013; 67: 56-60.
47. Piechota-Polańczyk A, Gorąca A. Influence of specific endothelin-1 receptor blockers on hemodynamic parameters and antioxidant status of plasma in LPS-induced endotoxemia. *Pharmacol Rep* 2012; 64: 1434-1441. doi: 10.1016/s1734-1140(12)70941-6.
48. Lorente L, Martín MM, Almeida T, et al. Association between serum total antioxidant capacity and mortality in severe septic patients. *J Crit Care* 2015; 30: 217.e7-217.e12. doi: 10.1016/j.jcrc.2014.09.012.
49. Martins PS, Brunialti MKC, Martos LSW, et al. Expression of cell surface receptors and oxidative metabolism modulation in the clinical continuum of sepsis. *Crit Care* 2008; 12: R25. doi: 10.1186/cc6801.
50. Rademann P, Weidinger A, Drechsler S, et al. Mitochondria-targeted antioxidants SkQ1 and MitoTEMPO failed to exert a long-term beneficial effect in murine polymicrobial sepsis. *Oxid Med Cell Longev* 2017; 2017: 6412682. doi: 10.1155/2017/6412682.
51. Cui W, Hu G, Peng J, Mu L, Liu J, Qiao L. Quercetin exerted protective effects in a rat model of sepsis via inhibition of reactive oxygen species (ROS) and downregulation of high mobility group box 1 (HMGB1) protein expression. *Med Sci Monit* 2019; 25: 5795-5800.
52. Barreiro E, Sánchez D, Gáldiz JB, Hussain SNA, Gea J; ENIGMA in COPD project. N-acetylcysteine increases manganese superoxide dismutase activity in septic rat diaphragms. *Oxid Med Cell Longev* 2005; 26: 1032-1039. doi: 10.1183/09031936.05.00003705.
53. Bergamini S, Rota C, Canali R, et al. N-acetylcysteine inhibits in vivo nitric oxide production by inducible nitric oxide synthase. *Nitric Oxide* 2001; 5: 349-360. doi: 10.1006/niox.2001.0356.
54. Ritter C, Andrades ME, Reinke A, Menna-Barreto S, Moreira JCF, Dal-Pizzol F. Treatment with N-acetylcysteine plus deferoxamine protects rats against oxidative stress and improves survival in sepsis. *Crit Care Med* 2004; 32: 342-349. doi: 10.1097/01.CCM.0000109454.13145.CA.
55. Wischmeyer PE, Dhaliwal R, McCall M, Ziegler TR, Heyland DK. Parenteral glutamine supplementation in critical illness: a systematic review. *Crit Care* 2014; 18: R76. doi: 10.1186/cc13836.
56. Buter H, Bakker AJ, Kingma WP, Koopmans M, Boerma EC. Plasma glutamine levels in patients after non-elective or elective ICU admission: an observational study. *BMC Anesthesiol* 2016; 16: 15. doi: 10.1186/s12871-016-0180-7.
57. Schaller G, Pleiner J, Mittermayer F, Posch M, Kapiotis S, Wolzt M. Effects of N-acetylcysteine against systemic and renal hemodynamic effects of endotoxin in healthy humans. *Crit Care Med* 2007; 35: 1869-1875. doi: 10.1097/01.CCM.0000275385.45557.25.
58. Rank N, Michel C, Haertel C, et al. N-acetylcysteine increases liver blood flow and improves liver function in septic shock patients: Results of a prospective, randomized, double-blind study. *Crit Care Med* 2000; 28: 3799-3807. doi: 10.1097/00003246-200012000-00006.
59. Prauchner CA. Oxidative stress in sepsis: Pathophysiological implications justifying antioxidant co-therapy. *Burns* 2017; 43: 471-485. doi: 10.1016/j.burns.2016.09.023.
60. Bar-Or D, Bar-Or R, Rael LT, Brody EN. Oxidative stress in severe acute illness. *Redox Biol* 2015; 4: 340-345. doi: 10.1016/j.redox.2015.01.006.
61. Andrews PJD, Avenell A, Noble DW, et al. Randomised trial of glutamine, selenium, or both, to supplement parenteral nutrition for criti-

- call ill patients. *BMJ* 2011; 342 (PG-d1542): d1542. doi: 10.1136/bmj.d1542.
62. Grintescu IM, Luca Vasiliu I, Cucereanu Badica I, et al. The influence of parenteral glutamine supplementation on glucose homeostasis in critically ill polytrauma patients – a randomized-controlled clinical study. *Clin Nutr* 2015; 34: 377-382. doi: 10.1016/j.clnu.2014.05.006.
 63. van Zanten a. RH, Hofman Z, Heyland DK. Consequences of the RE-DOXS and METAPLUS Trials: The End of an Era of Glutamine and Antioxidant Supplementation for Critically Ill Patients? *JPEN Parenter Enter Nutr* 2015; 39: 2014-2016. doi: 10.1177/0148607114567201.
 64. Heyland D, Muscedere J, Wischmeyer PE, et al. A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013; 368: 1489-1497. doi: 10.1056/NEJMoa1212722.
 65. Mokhtari V, Afsharian P, Shahhoseini M, Kalantar SM, Moini A. A review on various uses of n-acetyl cysteine. *Cell J* 2017; 19: 11-17. doi: 10.22074/cellj.2016.4872.
 66. Yazdi AP, Razavi M, Sheikh S, Boroumand N, Salehi M, Hashemy SI. Clinical trial assessment of intermittent and continuous infusion dose of N-acetylcysteine on redox status of the body in patients with sepsis admitted to the ICU. *J Intensive Care Med* 2020; 35: 1383-1388. doi: 10.1177/0885066618823152.
 67. Dodd S, Dean O, Copolov DL, Malhi GS, Berk M. Acetylcysteine for antioxidant therapy: pharmacology and clinical utility. *Expert Opin Biol Ther* 2008; 8: 1955-1962. doi: 10.1517/14728220802517901.
 68. Aldemir M, Koca HB, Bakı ED, et al. Effects of N-acetyl cysteine on renal functions evaluated by blood neutrophil gelatinase-associated lipocalin levels in geriatric patients undergoing coronary artery bypass grafting. *Anatol J Cardiol* 2016; 16: 504-511. doi: 10.5152/AnatolJCardiol.2015.6287.
 69. Dejager L, Pinheiro I, Dejonckheere E, Libert C. Cecal ligation and puncture : the gold standard model for polymicrobial sepsis? *Trends Microbiol* 2011; 19: 198-208. doi: 10.1016/j.tim.2011.01.001.
 70. Dyson A, Singer M. Animal models of sepsis: why does preclinical efficacy fail to translate to the clinical setting? *Crit Care Med* 2009; 37 (1 Suppl): S30-37. doi: 10.1097/CCM.0b013e3181922bd3.
 71. Esmon CT. Why do animal models (sometimes) fail to mimic human sepsis? *Crit Care Med* 2004; 32: 219-222. doi: 10.1097/01.ccm.0000127036.27343.48.
 72. Flores-Ruiz E, Miranda-Navales MG, Villasis-Keever MÁ. The research protocol VI : how to choose the appropriate statistical test. *Inferential statistics. Rev Alerg Mex* 2017; 64: 364-370.
 73. Amrhein V, Mcshane B. Retire statistical significance. *Nature* 2019; 567: 305-307. doi: 10.1038/d41586-019-00857-9.
 74. Lorente L, Martín MM, Ferreres J, et al. Serum caspase 3 levels are associated with early mortality in severe septic patients. *J Crit Care* 2016; 34: 103-106. doi: 10.1016/j.jcrc.2016.04.008.
 75. Lorente L, Martín MM, Gonzalez-Rivero AF, et al. Serum levels of caspase-cleaved cytokeratin-18 and mortality are associated in severe septic patients: pilot study. *PLoS One* 2014; 9: e109618. doi: 10.1371/journal.pone.0109618.
 76. Lorente L, Martín MM, González-Rivero AF, et al. Serum levels of caspase-cleaved cytokeratin-18 in patients with severe traumatic brain injury are associated with mortality: a pilot study. *PLoS One* 2015; 10: e0121739. doi: 10.1371/journal.pone.0121739